



CASE REPORT

Severe ANCA Associated Vasculitis in the Setting of Cocaine Abuse: A Case Report

Ana Ramirez Berlioz*, Orlando Garner, Eliana Wiesner, Alfredo Iardino Scarano and Kalpana Bhairavarasu

Internal Medicine, Texas Tech University Health Sciences Center, USA

*Corresponding author: Ana Ramirez Berlioz, PGY-3, Internal Medicine, Texas Tech University Health Sciences Center, USA, E-mail: ana.ramirez@ttuhsc.edu

Abstract

Introduction: Cocaine is a drug that is abused by 5 million Americans and most of the cocaine that comes to the United States has been tainted with Levamisole, a veterinary anti-helminthic, which recently has been linked to ANCA Associated Vasculitis (AAV) [1]. AAV is an unusual necrotizing vasculitis that attacks small vessels more often, and can have an ominous prognosis if left untreated [2].

Case description: We present a case of new onset P-ANCA associated vasculitis in a patient with a history of cocaine abuse who was lost to follow up and returned six years later with shortness of breath due to diffuse alveolar hemorrhage and heart failure with reduced ejection fraction, in addition to gross hematuria secondary to focal segmental necrotizing and sclerosing glomerulitis diagnosed with kidney biopsy in the setting of continued cocaine use.

The patient reappeared with purpuric lesions on her face, upper and lower extremities bilaterally with blebs and bullae that ulcerated. She was discharged after debridement but returned with worsened lesions when she discontinued wound care. Patient was readmitted with necrotic lesions complicated by myiasis; underwent re-debridement and was sent to a burn unit where she was treated with skin grafts.

Discussion: To our knowledge there are only five other cases with greater Total Body Surface Area (TBSA) involvement, only one other case that reported membranous nephropathy and no other report with full-thickness skin necrosis and lupus-like nephritis combining membranous and focal proliferative/sclerosing glomerulitis.

Since most of the cocaine in the US is adulterated with Levamisole, Levamisole induced AAV should be included as a differential diagnosis in patients presenting with characteristic purpuric lesions, hematuria, positive ANCA serology and history of cocaine abuse.

Case Report

A 41-year-old Hispanic female with known medical history significant for bronchial asthma presents to the Emergency Department (ED) in February 2010, complaining of a 3-week long history of an erythematous tender rash with irregular margins on the tip of her nose, upper and lower extremities (Figure 1) accompanied by bilateral elbow, knee and ankle pain. The rest of her physical exam and laboratory data was essentially normal, except for a Urine Drug Screen (UDS) which was positive for cocaine. She was managed symptomatically with antihistamines and steroid therapy with moderate improvement. Given presenting complaint of joint pain, she was worked up for possible connective tissue disease; ANA was positive, ANCA titers were 1:160, with positive Myeloperoxidase (MPO-ANCA), negative Proteinase 3 (PR3-ANCA) and a noted perinuclear staining pattern (P-ANCA). Anti-dsDNA antibodies, rheumatoid factor, and anti-smooth muscle antibodies were negative. (Please see Table 1 for detailed laboratory results). A skin biopsy was done which later demonstrated a neutrophil-rich infiltrate around the vessels of the dermis with fibrin thrombi within the superficial vessels. She was discharged with an oral prednisone taper but lost to follow-up.

Patient reappeared approximately 6 years later; when she complained of gross hematuria, chest pain, hemoptysis, and generalized weakness. Physical exam showed her to be pale, moderately hypertensive with a Blood Pressure (BP) of 159/82 mmHg, with fine rales



Figure 1: Skin lesions on initial presentation 2010.

Table 1: Lab work from initial encounter 2010.

Laboratory test	Date: 02/15/2010
WBC (Ref 4.5-10.0 k/UL)	4.5
Hemoglobin (Ref 12-16 g/dL)	14.4
Creatinine (Ref 0.6-1.2 mg/dL)	0.7
BUN (Ref 7-20.1 mg/dL)	11
Proteinuria	Negative
APTT (21.4-33.0)	37.0
PT (11.2-14.0)	13.0
INR	1.05
Myeloperoxidase Ab, IgG (Ref 0-19 Au/ml)	49
Serine Protease 3 Ab, IgG (Ref 0-19 Au/ml)	4
Antineutrophil Cytoplasmic Ab, IgG (Ref < 1:20)	1:160 **p-ANCA staining pattern observed and confirmed
DNA DS Ab (IU/ml) (Ref ≤ 4: negative 5-9: indeterminate ≥ 10: positive)	4
Actin (Smooth-muscle) Ab, IgG < 20: negative 20-30: weak positive > 30: high positive	< 20
Anti-nuclear antibody, IgG	Detected
ANA titer < 1:40 negative 1:40-1:80 Low antibody level > 1:80 Elevated antibody level	1:160 Speckled pattern

** : p-ANCA.

throughout the entirety of bilateral chest on auscultation. Review of initial laboratory data proved her to be leukopenic and anemic, with White Blood Cell (WBC) count of 3,200/uL, Hemoglobin (Hb) of 5.1 g/dL and

Hematocrit (HTC) of 15.6%; she was now also in acute renal failure with creatinine of 6.3 mg/dL and BUN 53 mg/dL. Her Brain Natriuretic Peptide (BNP) was elevated at 1963 pg/ml and a urinalysis was positive for protein (100 mg/dL), WBC's (14/hpf) and Red Blood Cells (RBC) (> 180/hpf). UDS was again positive for cocaine. A Computerized Tomography (CT) of the chest showed prominent bilateral patchy infiltrates and symmetric moderate pleural and small pericardial effusions (Figure 2); whereas abdominal CT scan was unremarkable except for mild pelvic fluid and mild soft tissue edema. Patient was admitted, transfused Packed Red Blood Cells (PRBC's), and started on high-dose steroid therapy. The work up done revealed diffuse alveolar hemorrhage diagnosed by bronchoscopy, acute kidney injury suspected to be due to rapidly progressive glomerulonephritis, anemia of chronic inflammation exacerbated by hematuria and new onset heart failure with grade III diastolic dysfunction and an ejection fraction of 45-50%. Immune serology continued to be P-ANCA positive, MPO-ANCA positive, and PR3-ANCA negative. (Please refer to Table 2 for details).

A kidney biopsy was done with preliminary report suggestive of type II/IV lupus nephritis/mixed connective tissue disease and final report showing focal segmental necrotizing and sclerosing glomerulitis (Figure 3).

Patient was diagnosed with AAV and discharged two weeks later with oral steroids, scheduled rituximab infusions, and instructions to follow up with Rheumatology and Nephrology specialists. She was seen shortly after as an outpatient, and her rituximab infusions were discontinued and switched to cyclophosphamide due to worsening renal function.

A few months later, patient was readmitted to the hospital for violaceous rash with bullae and blebs that became ulcers on face, upper and lower extremities bilaterally (Figure 4). Initial impression was a drug re-




Figure 2: CT scan thorax.

Table 2: Lab work from subsequent encounter 2016.

Laboratory test	01/25/16	01/26/16
Myeloperoxidase Ab, IgG (Ref \leq 19: negative 20-25: equivocal \geq 26: positive)	228	214
Serine Protease 3 Ab, IgG (Ref \leq 19: negative 20-25: equivocal \geq 26: positive)	9	5
Antineutrophil Cytoplasmic Ab, IgG (Ref < 1:20)		1:5120 **p-ANCA staining pattern observed and confirmed
DNA DS Ab (IU/ml) (Ref \leq 4: negative 5-9: indeterminate \geq 10: positive)	1:160 DS DNA IgG by IFA Detected by ELISA	
Anti-nuclear antibody, IgG ANA titer	Detected	Detected
< 1:40 negative 1:40-1:80 Low antibody level > 1:80 Elevated antibody level		1:160 Speckled pattern
Cyclic Citrullinated Peptide CCP AB, IgG \leq 19 Negative 20-39 Weak positive 40-59 Moderate positive \geq 60 Strong positive	4	
Cardiolipin Ab, IgG 0-14: Negative 15-19: Indeterminate 20-80: Low to moderately positive \geq 81 High positive		2
Cardiolipin Ab, IgM 0-12: Negative 13-19: Indeterminate 20-80: Low to moderately positive \geq 81 High positive		0

**: p-ANCA.

action versus infectious disease. Methylprednisolone was started, along with daptomycin and piperacillin/tazobactam, and a skin biopsy was performed which showed minimal inflammation with no signs of leukocytic vasculitis. Vasculitis workup including CPK, aldolase, dsDNA, C3, C4, ANCA and Beta-2 glycoprotein came back negative and wound culture grew coagulase negative *S. aureus*. Antibiotic therapy was deescalated to daptomycin, an eschar was beginning to form and patient was discharged three weeks later, on prednisone, amoxicillin/clavulanate, doxycycline, and cyclophosphamide. Patient was seen as an outpatient one week af-




Figure 3: Renal biopsy electron microscopy, mid-magnification.



Figure 4: Skin lesions on subsequent presentation in 2016, demonstrating violaceous rash with bullae, blebs and ulceration.



Figure 5: Infected skin lesions on readmission in 2016, with necrosis and myiasis.

ter discharge where she stated she was doing her own wound care and having no complications on prednisone and cyclophosphamide, wounds were still healing.

Three months later patient was readmitted because the ulcers in her upper and lower extremities became necrotic complicated with foul smell and wound myiasis after she discontinued wound care (Figure 5). Physical examination on admission showed full-thickness loss with granulating and necrotic tissue, accompanied by visible larvae in upper and lower extremities bilaterally. Antibiotic therapy was started with daptomycin and ceftazolin; and surgical debridement was performed. Due to the extent of the debridement she was transferred to an institution with a burn unit for higher level of care and skin grafting was performed.

She was seen again three months later as an outpatient after skin grafts were done, wounds were healing well, no amputations were performed, renal function was stable and currently patient is only receiving prednisone with no evident complications.

Discussion

Levamisole is an anti-helminthic, historically used in humans to treat cancer and rheumatologic diseases, given its immunomodulating properties. It has been banned by the Food and Drug Administration (FDA) for use in humans since 2000, due to its side effects of agranulocytosis and vasculitis. However, it continues to be available for veterinary use and is a known adulterant for cocaine, as it adds bulk and weight, potentially intensifying cocaine's psychotropic effects [3].

Levamisole induced AAV is a relatively new entity with cases first being documented as early as 2009; its clinical presentation being a tender, purpuric rash in extremities, trunk, digits and nasal tip [4]. Considering that in 2016, 93% of the cocaine seized by the US Drug Enforcement Agency was cut with Levamisole and/or levamisole mixtures [5], and that the patient's initial clinical presentation was similar to other cases described in literature; in retrospect, levamisole induced AAV is the likely culprit for this patient's initial symptoms in 2010.

The presence of Levamisole cannot be recognized by a simple UDS. Exposure to the substance can be con-

firmed with Gas Chromatography-Mass Spectrometry (GC-MS) in urine [6] but this is not a readily available test in most hospitals. It may be requested in patients who present with similar rash, positive ANCA and history of cocaine abuse. However, according to some authors the identification of the substance may not be necessary to make the diagnosis [3].

The natural history of the cutaneous manifestations of levamisole induced AAV is spontaneous resolution after the substance is withdrawn; though immunologic markers may persist for up to 14 months [7]. In cocaine-induced systemic vasculitis, patients may be positive for PR3-ANCA, MPO-ANCA and/or the combination PR3-ANCA and MPO-ANCA may be found [8].

To our knowledge there are no defined criteria for the identification of levamisole induced AAV and its diagnosis remains based on exclusion. In recent years, some authors have proposed that once diagnosis is suspected, the staging of this condition be based on the clinical cutaneous findings, as such: The mild stage (I) consists of livedo with mild purpura and no areas of necrosis. The moderate stage (II) has fixed purpuric patches and or plaques, without significant skin ulceration and the severe stage (III) is characterized by extensi vereticular (IIIa) or confluent (IIIb) purpura accompanied by bullous involvement with or without cutaneous necrosis [9].

Although there are currently no definitive guidelines on the treatment of levamisole induced vasculitis, besides avoiding further exposure to the substance, the management is essentially supportive [1]. To our knowledge, four cases of severe levamisole-induced AAV have been described with extensive full-thickness skin necrosis, involving 15, 52, 49 and ~20% of TBSA [10,11]. On these patients, invasive therapy was performed including surgical debridement, skin grafting and amputations, with variable outcomes. A recently published article demonstrated that early excision and grafting of large surface levamisole-induced skin necrosis in dependent of autoantibodies results in successful outcomes and limits wound morbidity [10]. Our patient's TBSA involvement was estimated at 13% and she eventually underwent extensive skin grafting and was managed in a burn unit, with good end results.

Renal biopsy findings were also remarkable since it reported a lupus-like nephritis combining membranous and focal proliferative/sclerosing glomerulitis which to our knowledge has not been reported in cases of Levamisole-induced AAV, even though there have been reports of two cases of membranous nephropathy with crescents [12,13].

Our patient's course was complicated because she was lost to follow up for six years and remained untreated, presumably still using cocaine, as inferred from her subsequent positive UDS. She was seen again with a wide constellation of problems and multi-system in-

volvement including diffuse alveolar hemorrhage and focal segmental necrotizing and sclerosing glomerulitis diagnosed by kidney biopsy. It is well known that patients with AAV have an increased mortality at one year if left untreated but the patient survived all these years without any therapeutic regimen [14].

Conclusion

The presence of a characteristic clinical picture and the presumed widespread use of levamisole as an adulterant for cocaine should raise suspicions for levamisole induced AAV whenever patients are seen with purpuric rashes, history of cocaine abuse and positive ANCA.

Remission of the disease occurs spontaneously after stopping the use of cocaine and patients should be strongly encouraged to abstain from it, since prolonged use may further cause systemic damage.

References

1. Patnaik S, Balderia P, Vanchhawng L, Markazi P, Wykretowicz J, et al. (2015) Is Levamisole-Induced Vasculitis a Religated Diagnostic Possibility? A Case Report and Review of Literature. *Am J Case Rep* 16: 658-662.
2. Houben E, Bax WA, van Dam B, Sliker WA, Verhave G, et al. (2016) Diagnosing ANCA-associated vasculitis in ANCA positive patients A retrospective analysis on the role of clinical symptoms and the ANCA titre. *Medicine* 95: e5096.
3. Larocque A, Hoffman R (2012) Levamisole in cocaine: Unexpected news from an old acquaintance. *Clin Toxicol* 50: 231-241.
4. Roberts JA, Chevez-Barríos P (2015) Levamisole-Induced Vasculitis A Characteristic Cutaneous Vasculitis Associated With Levamisole-Adulterated Cocaine. *Arch Pathol Lab Med* 139: 1058-1061.
5. US Department of Justice and Drug Enforcement Administration (2016) 2016 National Drug Threat Assessment Summary.
6. Lynch KL, Dominy SS, Graf J, Kral AH (2011) Detection of Levamisole Exposure in Cocaine Users by Liquid Chromatography-Tandem Mass Spectrometry. *J Anal Toxicol* 35: 176-178.
7. Gross RL, Brucker J, Bahce-Altuntas A, Abadi MA, Lipoff J, et al. (2011) A novel cutaneous vasculitis syndrome induced by levamisole-contaminated cocaine. *Clin Rheumatol* 30: 1385-1392.
8. Cohen Tervaert JW, Damoiseaux J (2012) Antineutrophil cytoplasmic autoantibodies: how are they detected and what is their use for diagnosis, classification and follow-up? *Clin Rev Allergy Immunol* 43: 211-219.
9. Gillis JA, Green P, Williams J (2014) Levamisole-induced vasculopathy: Staging and management. *J Plast Reconstr Aesthet Surg* 67: e29-e31.
10. Miner J, Gruber P, Perry TL (2015) Early excision and grafting, an alternative approach to the surgical management of large body surface area levamisole-adulterated cocaine induced skin necrosis. *Burns* 41: e34-e40.
11. Alekseyev K, Micaily I, Parikh N (2016) A Severe Case of Levamisole-Induced Vasculitis Requiring Extensive Surgery and Skin Grafts. *J Cutan Aesthet Surg* 9: 41-43.
12. Moinuddin I, Madhrira M, Bracamonte E, Thajudeen B, Sussman A (2016) Membranous nephropathy with crescents associated with levamisole-induced MPO-ANCA vasculitis. *Pathol Res Pract* 212: 650-653.
13. Roca-Argente L, Moll-Guillen JL, Espi-Reig J, Blanes-Julia M, García-Martínez AM, et al. (2015) Membranous glomerulonephritis an cellular crescents induced by levamisole-adulterated cocaine abuse: a case report. *Ann Transl Med* 3: 271.
14. Harris A, Szilveszter D, Polner K (2017) Therapy and prognosis of ANCA-associated vasculitis from the clinical nephrologist's perspective. *Int Urol Nephrol* 49: 91-102.