



## ORIGINAL ARTICLE

## Poverty, Urban Environmental Factors and Spatial Distribution of Visceral Leishmaniasis in an Area of Recent Emergence in Inner Brazil

José Cláudio Simão, PhD<sup>1\*</sup> , Cassiano Victória, PhD<sup>2</sup>  and Carlos Magno Castelo Branco Fortaleza, MD, PhD<sup>1</sup> 



<sup>1</sup>Department of Tropical Diseases, Botucatu Medical School, São Paulo State University (UNESP), São Paulo, Brazil

<sup>2</sup>Department of Veterinary Hygiene and Public Health, Faculty of Veterinary, São Paulo State University (UNESP), São Paulo, Brazil

\*Corresponding author: José Cláudio Simão, Department of Tropical Diseases, Botucatu Medical School, São Paulo State University (UNESP), Botucatu, São Paulo, Distrito de Rubião Junior, S/N CEP/ZIP 18618-970, Brazil, Tel: +55-14-3882-1284, Fax: +55-14-3811-1515

### Abstract

In an urban area of recent emergence of visceral leishmaniasis (VL) in Brazil, the spatial distribution of both VL and VL-HIV coinfection was negatively associated with per capita income, population density, paving of streets, existence of sidewalks and drains for rainfall. Poverty and urban infrastructure deficits may influence VL emergence.

### Keywords

Visceral leishmaniasis, Leishmania-HIV coinfection, Spatialepidemiology

### Introduction

In the past three decades, visceral leishmaniasis (VL) in Brazil changed from rural endemics of Northeastern States into an emergent disease of great urban centers [1].

The area affected spread southwards affecting São Paulo, the most populous state in the country. In this setting, the epidemics of VL and AIDS intertwined [2].

In urban areas of recent emergence, the spatial distribution of VL cases is not homogeneous [3]. Socio-economic and environmental factors may contribute to the occurrence of LV in specific areas [4].

Therefore, the study of those factors is likely to

provide clues to determinants of the emergence of VL.

### The Study

We studied the spatial distribution of overall VL and VL-HIV coinfection in the city of Bauru (340,000 inhabitants), inner São Paulo State, Brazil. The first notifications of VL in that city date from 2003. The cumulative incidence rates of VL and VL-HIV (per 100,000 inhabitants) in the period 2003-2016 were 131.1 and 19.7, respectively.

The addresses of cases were georeferenced in QGIS 2.18 (QGIS Development Team [2016]. QGIS Geographic Information System. Open Source Geospatial Foundation Project. <http://qgis.osgeo.org>), using UFT-8 codifications and MMQGIS geocodification algorithm. After georeferencing, cases were assigned to census sectors of the city of Bauru, Brazil, according to Brazilian Institute of Geography and Statistics (IBGE). Kernel density maps were generated in ArcGis 10.1 (ESRI, Redlands, CA) using the following algorithms: “integrate”, “collect events” and “hotspot analysis”.


Census sectors were the units for analysis of predictors of VL and VL-HIV incidence. Variables included in the analysis were collected in the 2010 Brazilian Census Data (IBGE) and included and per capita



**Citation:** Simão JC, Victória C, Fortaleza CMCB (2022) Poverty, Urban Environmental Factors and Spatial Distribution of Visceral Leishmaniasis in an Area of Recent Emergence in Inner Brazil. Int Arch Public Health Community Med 6:081. doi.org/10.23937/2643-4512/1710081

**Accepted:** June 30, 2022; **Published:** July 02, 2022

**Copyright:** © 2022 Simão JC, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.



income and population density. Other data collected from the same source included percentage of houses in places with sidewalks, paving of streets, drains for rainfall, trees, sewage and garbage in the open. Data were analyzed in Stata 14 (College Station, TX), using univariate and multivariable (single-step) models of Zero-inflated Poisson Regression.

Figure 1 presents spatial distribution of cases, in crude incidence per census sectors and Kernel densities. Predictors of geographic incidence are represented in Table 1. Briefly, overall VL incidence was negatively associated with population density and per capita income, while there was negative association of the incidence of VL-HIV coinfection with per capita income and presence of drains for rainfall.

Our results provide clues to the routes of urbanization

of VL and intersection of VL and AIDS epidemics. In univariable analysis, several aspects linked to poverty (e.g., absence of sidewalks and paving of streets, presence of garbage in the open) were associated with the outcomes. Those environmental factors may be proxies for both the presence of vectors (*Lutzomyia longipalpis*) breeding sites (usually sites with decaying organic matter) and of reservoirs (domesticated or stray dogs) [5,6].

Similarly to our results, proxy indicators of poverty such as (illiteracy and income) have been described both in areas of early [7] and recent emergence [8]. Urban VL has been associated with living in the periphery of cities, which are sites where favelas and poor neighborhoods are located. Not surprisingly, those are areas of greater prevalence of HIV infection [9].

**Table 1:** Factors associated with the incidence of Visceral Leishmaniasis (VL) and VL-HIV coinfection among census sectors in the city of Bauru, inner Brazil.

Factors	Univariada		Multivariada	
	IRR (95%CI)	P	IRR (95%CI)	P
<b>Overall VL</b>				
Population density	<b>0.95 (0.92-0.99)</b>	<b>0.006</b>	<b>0.94 (0.90-0.97)</b>	<b>&lt; 0.001</b>
Per capita income (BRL 100)	<b>0.90 (0.87-0.92)</b>	<b>&lt; 0.001</b>	<b>0.89 (0.86-0.92)</b>	<b>&lt; 0.001</b>
Street paving (x10%)	<b>0.93 (0.90-0.96)</b>	<b>&lt; 0.001</b>	1.01 (0.92-1.10)	0.87
Sidewalks (10%)	<b>0.92 (0.90-0.95)</b>	<b>&lt; 0.001</b>	0.99 (0.91-1.09)	0.91
Rain drains (10%)	<b>0.94 (0.90-0.98)</b>	<b>0.002</b>	0.96 (0.92-1.00)	0.08
Trees (x10%)	1.03 (0.92-1.10)	0.57	0.92 (0.80-1.05)	0.21
Open sewer (10%)	1.08 (0.93-1.27)	0.33	1.09 (0.91-1.31)	0.36
Open-air garbage (x10%)	<b>1.06 (1.02-1.12)</b>	<b>0.004</b>	1.00 (0.95-1.06)	0.96
<b>Coinfecção LV-HIV</b>				
Population density	0.97 (0.90-1.05)	0.48	0.94 (0.86-1.03)	0.18
Per capita income (R\$ 100))	<b>0.85 (0.78-0.93)</b>	<b>&lt; 0.001</b>	<b>0.80 (0.71-0.91)</b>	<b>&lt; 0.001</b>
Street paving (x10%)	<b>0.92 (0.85-0.98)</b>	<b>0.02</b>	0.95 (0.90-1.19)	0.67
Sidewalks (10%)	<b>0.92 (0.86-0.99)</b>	<b>0.03</b>	1.09 (0.87-1.37)	0.44
Rain drains (10%)	<b>0.86 (0.77-0.95)</b>	<b>0.004</b>	<b>0.87 (0.78-0.98)</b>	<b>0.02</b>
Trees (x10%)	1.04 (0.81-1.32)	0.76	0.94 (0.73-1.19)	0.60
Open sewer (10%)	11.08 (0.66-1.78)	0.76	1.27 (0.76-2.13)	0.36
Open sewer (10%)	0.99 (0.87-1.13)	0.89	0.86 (0.74-1.01)	0.07

VL: visceral leishmaniasis; HIV: Human Immunodeficiency Virus; BRL: Brazilian currency (Real, approximately US\$0.25); IRR: incidence rate; CI: confidence interval.

## Conclusions

Taken together, all those findings reinforce the social and economic determination in the distribution of VL and VL-coinfection. Since a recent systematic review failed to identify effective measures to prevent VL in Latin America [10], interventions aimed at improving income and housing conditions may be pathways for controlling this disease.

## References

- Arias JR, Monteiro PS, Zicker F (1996) The reemergence of visceral leishmaniasis in Brazil. *Emerg Infect Dis* 2: 145-146.
- Leite de Sousa-Gomes M, Romero GAS, Werneck GL (2017) Visceral leishmaniasis and HIV/AIDS in Brazil: Are we aware enough? *PLoS Negl Trop Dis* 11: e0005772.
- Cardim MF, Vieira CP, Chiaravalloti-Neto F (2015) Spatial and spatiotemporal occurrence of human visceral leishmaniasis in Adamantina, State of São Paulo, Brazil. *Rev Soc Bras Med Trop* 48: 716-723.
- Ursine RL, Dias JV, Morais HA, Pires HH (2016) Human and canine visceral leishmaniasis in an emerging focus in Araçuaí, Minas Gerais: spatial distribution and socio-environmental factors. *Mem Inst Oswaldo Cruz* 111: 505-511.
- Casanova C, Andrighetti MT, Sampaio SM, Marcoris ML, Colla-Jacques FE, et al. (2013) Larval breeding sites of *Lutzomyia longipalpis* (Diptera: Psychodidae) in visceral leishmaniasis endemic urban areas in Southeastern Brazil. *PLoS Negl Trop Dis* 7: e2443.
- Teixeira-Neto RG, da Silva ES, Nascimento RA, Belo VS, de Oliveira CD, et al. (2014) Canine visceral leishmaniasis in an urban setting of Southeastern Brazil: an ecological study involving spatial analysis. *Parasit Vectors* 7: 485.
- de Almeida AS, Medronho R de A, Werneck GL (2011) Identification of risk areas for visceral leishmaniasis in Teresina, Piauí State, Brazil. *Am J Trop Med Hyg* 84: 681-687.
- de Araújo VE, Pinheiro LC, Almeida MC, de Menezes FC, Morais MH, et al. (2013) Relative risk of visceral leishmaniasis in Brazil: a spatial analysis in urban area. *PLoS Negl Trop Dis* 7: e2540.
- Alves AT, Nobre FF, Waller LA (2016) Exploring spatial patterns in the associations between local AIDS incidence and socioeconomic and demographic variables in the state of Rio de Janeiro, Brazil. *Spat Spatiotemporal Epidemiol* 17: 85-93.
- Romero GA, Boelaert M (2010) Control of visceral leishmaniasis in Latin America - a systematic review. *PLoS Negl Trop Dis* 4: e584.