



CASE REPORT

Embolization in the Treatment of a Major Retropubic Hemorrhage Following Tension-Free Vaginal Tape: A Case Report and Literature Review

Margrethe Foss Hansen^{1*}, Mads Henrik Carstensen², Jens Prien-Larsen³ and Lars Alling-Møller⁴



¹Department of Obstetrics and Gynecology, Nordsjællands Hospital, Denmark

²Department of Gastrointestinal Surgery, Slagelse Hospital, Denmark

³Department of Obstetrics and Gynecology, Nykøbing Falster Hospital, Denmark

⁴Department of Obstetrics and Gynecology, Zealand University Hospital, Denmark; Department of Clinical Medicine, University of Copenhagen, Denmark

*Corresponding author: Margrethe Foss Hansen, (ORCID 0000-0002-0497-5070), Department of Obstetrics and Gynecology, Nordsjællands Hospital, Dyrehavevej 29, 3400, Hillerød, Denmark, Tel: +45-40427966, Fax: +45-482937, E-mail: margrethefoss@hotmail.com

Abstract

55-year-old para 2 woman was brought to the emergency department following insertion of a tension-free vaginal tape complaining of lower abdominal pain and vertigo. The subsequent angiography localized a significant bleeding from the right obturator artery and a retropubic hematoma 17.5 cm × 10.0 cm in size. The patient was successfully treated by arterial embolization; however, the postoperative course was greatly protracted due to persistent subfebrilia and eventually the hematoma was removed by laparoscopy. This case illustrates that embolization should be considered an important alternative for management of arterial bleeding following synthetic midurethral slings. However, removal by laparoscopy or laparotomy may be necessary in large hematomas subsequently to embolization.

Keywords

Arterial bleeding, Tension-Free vaginal tape, Retropubic hemorrhage, Embolization

Introduction

The Tension-Free Vaginal Tape (TVT) was introduced in 1996 and is now widely used in surgical treatment of urinary incontinence due its low risk of surgical complications and high long-term cure rate [1]. Surgical complications may however occur as the blind passage of a

sharp instrument through the retropubic space carries the risk of visceral or vascular injury [1,2]. The Transobturator Tapes (TVT-O, TOT) were introduced later to minimize the risk of TVT. However, vascular and nerve injuries due to these procedures have also been reported [1].

Venous injuries resulting in a vulval-vaginal or retropubic hematoma are not uncommon [1,3]. An arterial injury is rare but potentially life-threatening due to the risk of blood loss and hemodynamic instability. Commonly, it has been managed through an open abdominal incision [1,3]. In the last decade; embolization has gained favor for the treatment of acute hemorrhage in different anatomical areas within several surgical specialisms for the treatment of acute hemorrhage [4]. Case reports have shown favorable outcomes of arterial embolization following insertion of a synthetic Midurethral Sling (MUS) complicated by arterial hemorrhage [5-13].

This case report describes an injury to the right obturator artery following a TVT procedure complicated by arterial hemorrhage successfully treated by embolization, but with a protracted postoperative course. The




Figure 1: Pelvic arteriography before embolization of the right obturator artery.




Figure 2: Enhanced computed tomography shows a major retropubic hematoma (17.5 cm × 10 cm) following tension-free vaginal tape.

present case is discussed with reference to a brief systematic review of the literature.

Case Report

Seven hours after undergoing an uneventful TVT insertion under local anesthesia a 55-year-old non-obese para 2 woman was brought to the emergency department complaining of lower abdominal pain and vertigo. On physical examination, she was alert and aware, but pale, peripherally cold and hypotensive (blood pressure was 80/30 and pulse = 55/minute). Apart from medical treatment for hypertension, she was healthy. Prior surgical history included an appendectomy only. On suspicion of intraperitoneal bleeding, laparoscopic evaluation was performed showing a large hematoma retropubically and on the posterior abdominal wall and after conferral with a vascular surgeon, the decision was made to perform arterial embolization. The subsequent angiography localized significant bleeding from the right obturator artery; an anterior branch of the internal iliac vessel (Figure 1) and a retropubic hematoma 17.5 cm × 10.0 cm in size (Figure 2). The patient was treated by selective embolization with coils of the right obturator artery (Figure 3) and her vital signs remained stable after the procedure. Five days postoperatively, the patients presented with fever of 38.9 °C and diffuse lower abdominal pain. She received intravenous broad-spectrum antibiotics, the pain abated and an enhanced Computed Tomography (CT) scan revealed a reduced hematoma sized 11 cm × 6 cm. To prevent formation of an abscess and due to ongoing fever (temperature up to 38.9 °C) laparoscopic evacuation of the hematoma was performed with a total blood loss of 1000 ml on postoperative day 10. Because the upper part of the tape had loosened from the tissue and was visible intraperitoneally, this part had to be removed.

The patient was in fair general health postoperative-



Figure 3: Pelvic arteriography after embolization of the right obturator artery.

ly, but due to low-grade fever, she received intravenous antibiotics for nine days in total. On postoperative day 14, the patient was discharged on oral antibiotics. Due to persistent low-grade fever on postoperative day 21, an enhanced CT scan was performed excluding injury to the urinary tract, however showing a 1.4 cm × 6 cm pelvic mass representing either a small abscess or a residual hematoma. No further treatment was initiated. Two months postoperatively, the patient was well-being and reported no stress urinary incontinence.

Discussion

To decrease the risk of vascular injury it is imperative that the surgeon employs the correct technique [6,14] and ensures there no bleeding occurs from the vaginal incisions perioperatively [1]. However, since it is not uncommon for the pelvic arteries to follow an aberrant course [14], hemorrhage is not always possible to prevent. Rarely, hemorrhage results from an occult bleeding diathesis [6]. In the present case, while there was no explanation for the hemorrhage, as it was unlikely to

have been due to surgical technique, the passage of the trocars through the retropubic space was inevitably a blind procedure [1] and vessel damage was hence more likely to have been a chance adverse event.

The incidence of retropubic hematomas following MUS sling procedure using TVT has been reported as 2 - 4% and the consequent need for surgical intervention as 0.3 - 1.2% [2].

The majority of retropubic hematomas is due to venous bleeding and can often be managed expectantly [1]. Arterial bleeding, however, is a rare event and even though immediate and adequate treatment is imperative, no standard for its management has been established. Commonly, it has been managed through an open abdominal incision [1,3].

To date, eleven cases have described arterial bleeding following synthetic MUS managed by arterial embolization [5-13] (Table 1). The present case, in agreement with the other studies, illustrates the usefulness of angiography to locate the bleeding and to obtain hemosta-

sis in arterial injuries following synthetic MUSs [5-13]. Unlike previous cases, the present postoperative course was greatly protracted due to persistent infection, multiple assessments and ultimately, removal of the hematoma. There are several further possible explanations for the protracted course: Firstly, unlike in the majority of reported cases, the patient underwent laparoscopy prior to angiography [5,7-12] which may have increased the risk of infection. Secondly, an obvious disadvantage of leaving the hematoma *in situ* is that the stagnant blood provides an ideal environment for bacterial infection to develop [1]. The risk in the present case might conceivably have been reduced had the patient been given prophylactic antibiotics, as recommended in some earlier case reports [1,3].

Thirdly, the size of the hematoma might also be a limiting factor, as this is the largest hematoma (10 cm × 17.5 cm) reported following a synthetic MUS, for which management has been attempted by embolization without subsequent removal [6-8,12,13]. In the literature, hematomas up to 13 - 16 cm following synthetic MUSs have been

Table 1: Literature on arterial injury following synthetic midurethral slings and embolization.

Case (°)	Author	Symptoms presented (hours/days)	Presenting symptoms and signs	Arterial injury	Size of hematoma	Blood transfusion	Assessment	Management	Follow-up
1	Elard, et al. [5]	Two days	Anemia	Inferior vesical artery	-	Yes	CT, angiography	Embolization	Postoperative course uneventful
2	Zorn KC [6]	Six hours	Bradycardia Hypovolemia	Obturator artery	13.9 cm × 8.5 cm	Yes 4 units of packed red blood cells	Laparoscopic evaluation angiography	Embolization (coils)	Postoperative course uneventful Improvement in stress incontinence
3	Takatera H, et al. [7]	Three hours	Lower abdominal pain, abdominal distension and nausea	-	16 cm × 12 cm × 11 cm	No	CT, angiography	Embolization (gel foam, coils)	Postoperative course uneventful No surgical removal of hematoma
4	Jung, et al. [8]	One hour	Lower abdominal pain, abdominal distension and nausea Bradycardia	Internal pudendal artery	10 cm × 9 cm × 8 cm	No	CT, angiography	Embolization (gel foam)	Postoperative course uneventful Two years follow-up: No stress urinary incontinence Mild urge incontinence
5	Huffaker RK [9]	Two hours	Hypotensive Anemia, Palpable supra-pubic mass	Obturator artery+collaterals	-	Yes 2 units of packed red blood cells	Angiography	Embolization (coils + gel foam)	Discharge on day 2 with Foley catheter. Feeling well one week postoperatively
6	Ku CH [10]	-	-	-	-	-	-	-	-

7	Krynytska I [11]	Two hours	Severe vulval pain	Obturator artery	-	Yes 2 units of packed red blood cells	Angiography	Embolization (coils) Subsequent evacuation of hematoma, general anaesthesia (Reduce pain, infection)	Folley catheter removed on day 10 Tenderness up to six months No stress urinary incontinence, bladder overactivity or voiding dysfunction
8	Ko JK [12]	One day	Perineal pain, Oozing from vagina	Internal pudendal artery	7 cm	Yes 4 units of packed red blood cells	CT, angiography	Embolization (coils + gel foam)	Discharge on day 3. Vaginal spotting and pain on day 7, edematous and fragile tissue on anterior vaginal wall. No significant bleeding. Wound completely healed at day 21. No further symptoms.
9	Cho EJ [13] Case A	One day	Abdominal pain Voiding difficulty	Upper vaginal branch of iliac int.	-	Yes 2 units of packed red blood cells	Laparotomy Angiography	Removal of hematoma laparotomy Unknown bleeding focus Subsequent embolization (gel foam)	Discharge on day 4 Presentation on day 12 with lower abdominal pain. Computed tomography-reduction of hematoma Improvement of stress incontinence on three months follow up
	Case B	Perioperatively	Lower abdominal pain	Inferior epigastric artery	10 cm	Yes 8 units of packed red blood cells	CT, Laparotomy Angiography	Removal of hematoma laparotomy Unknown bleeding focus Subsequent embolization (microcoil + gel foam)	Discharge on day 5 Presentation on day 7 with lower abdominal pain. Computed tomography showed a size of residual hematoma of 7 cm. On day 13, no further symptoms
	Case C	Intraoperatively	Intraoperative bleeding from Incision in suburethra	Anterior trunk of iliac artery		Yes 8 units of packed red blood cells + 4 fresh frozen plasma	Angiography	Embolization (gel foam)	Discharge and Folley catheter removed on day 7 without any complications.

(*) Case 1-5, 9(C) Embolization following insertion of a TVT. Case 6-9 (A,B) Embolization following insertion of a TOT.

managed with embolization without the need for subsequent removal [6,7]. Normally, small hematomas (< 5 cm) do not require any surgical intervention [1], but hematomas up to 10 cm have in stable patients been managed expectantly [3], which suggests that the maximum size for embolization also lies within this range.

The present case report, in agreement with existing literature, illustrates that embolization should be considered an important alternative treatment for management of arterial bleeding following synthetic MUSs; as the minimal approach of the TVT is preserved while managing a potentially life-threatening condition. However, the size of the hematoma might be a limiting factor for leaving it *in situ*, which necessitates removal by laparoscopy or laparotomy subsequently to embolization.

Lack of access to intervention radiology and awareness of the treatment among uro-gynecologists may reduce this surgical practice.

Conflict of Interest

No potential conflict of interest relevant to this article was reported.

Grant/Fund Support

None.

Research Ethics

Written informed content was obtained from the patient for publication of this case and any accompanying images.

References

- Balachandran A, Curtiss N, Duckett J (2015) The management of massive haematomas after insertion of retropubic mid-urethral slings. *Int Urogynecol J* 26: 1449-1452.
- Flock F, Reich A, Muche R, Kreienberg R, Reister F (2004) Hemorrhagic complications associated with tension-free vaginal tape procedure. *Obstet Gynecol* 104: 989-994.
- Kölle D, Tamussino K, Hanzal E, Tammaa A, Preyer O, et al. (2005) Bleeding complications with the tension-free vaginal tape operation. *Am J Obstet Gynecol* 193: 2045-2049.
- Velmahos GC, Toutouzas KG, Vassiliu P, Sarkisyan G, Chan LS, et al. (2002) A prospective study on the safety and efficacy of angiographic embolization for pelvic and visceral injuries. *J Trauma* 53: 303-308.
- Elard S, Cicco A, Salomon L, You K, de La Taille A, et al. (2002) Embolization for arterial injury incurred during tension-free vaginal tape procedure. *J Urol* 168: 1503.
- Zorn KC, Daigle S, Belzile F, Tu le M (2005) Embolization of a massive retropubic hemorrhage following a tension-free vaginal tape (TVT) procedure: case report and literature review. *Can J Urol* 12: 2560-2563.
- Takatera H, Mukai M, Fujii T, Nishi M, Okada H, et al. (2006) Arterial injury associated with tension-free vaginal tape procedure successfully treated by radiological embolization: a case report. *Hinyokika Kiyo* 52: 741-743.
- Jung YS, Lee JH, Shin TS, Han CH, Kang SH, et al. (2010) Arterial Injury Associated with Tension-Free Vaginal Tapes-SECUR Procedure Successfully Treated by Radiological Embolization. *Int Neurourol J* 14: 275-277.
- Huffaker RK, Handcock T, Muir TW (2010) Hemorrhagic complication during placement of tension-free vaginal tape. *Female Pelvic Med Reconstr Surg* 16: 365-367.
- Chun Hoe Ku, Ji Sung Lee, Jeong Ho Kim (2009) A case using artery embolization for the patient with uterine bleeding after transobturator tape operation. *J Korean Continence Soc* 13: 173-176.
- Krynytska I, Holmes R (2013) Vulval haematoma after TVT-obturator insertion requiring arterial embolisation. *J Obstet Gynaecol* 33: 908-909.
- Ko JK, Ku CH (2014) Embolization for pelvic arterial bleeding following a transobturator tape procedure. *J Obstet Gynaecol Res* 40: 865-868.
- Cho EJ, Kim JB, Park SY, Kim SH, Kim CH, et al. (2016) Pelvic artery embolization in the management of pelvic arterial bleeding following midurethral sling surgery for stress urinary incontinence. *Obstet Gynecol Sci* 59: 163-167.
- Muir TW, Tulikangas PK, Fidela Paraiso M, Walters MD (2003) The relationship of tension-free vaginal tape insertion and the vascular anatomy. *Obstet Gynecol* 101: 933-936.