



RESEARCH ARTICLE

Can Pre-Operative Endoscopy Identify Patients at Risk for Gastroesophageal Reflux Disease after Sleeve Gastrectomy?

Rami R Mustafa^{1,4*}, Leena Khaitan¹, Michał Robert Janik^{1,5}, Adel Alhaj Saleh^{1,3}, Mohammed Alshehri¹, Seyed Mohammad Kalantar Motamedi¹, Adil Khan¹, Heba elghalban², Tomasz Rogula¹ and Mujjahid Abbas¹

¹Bariatric Division, Department of Surgery, University Hospitals, Cleveland Medical Center/Case Western Reserve University, Cleveland, Ohio, USA

²Department of Family Medicine, University Hospitals, Cleveland Medical Center/Case Western Reserve University, Cleveland, Ohio, USA

³Department of Surgery, Texas Tech University Health Sciences Center, Lubbock TX, USA

⁴Department of Surgery, University Hospitals, Menoufia/Faculty of Medicine, Menoufia, Egypt

⁵Department of General, Oncologic, Metabolic and Thoracic Surgery, Military Institute of Medicine, Warszawa, Poland

*Corresponding author: Rami Mustafa, MD, Bariatric Division, Department of Surgery, University Hospitals, Cleveland Medical Center/Case Western Reserve University, 11100 Euclid Ave, Cleveland, Ohio 44121, USA, E-mail: rami.rabee@med.menoufia.edu.eg



Abstract

Background: Obese individuals have higher incidence of hiatal hernia (HH). There is controversy over the need for preoperative esophagogastroduodenoscopy (EGD) before bariatric procedures. The aim of this study is to determine the predictive value of preoperative endoscopy in diagnosing HH.

Methods: Under IRB Approval A retrospective review of 402 cases who underwent sleeve gastrectomy (SG) in our academic center between January 2011 and December 2015 was performed. Patients who had preoperative EGD reports were enrolled in the study. Data were collected for the demographics, preoperative endoscopic and intraoperative findings. Each patient's EGD findings were compared with the intraoperative findings.

Results: From total of 402 SG cases, 381 (81% female) had preoperative EGD. The mean age of the subjects was 45.8 years (\pm 10.6 years) with the mean BMI of 47.5 kg/m² (\pm 8.8 kg/m²). There was no significant difference in age and BMI between males and females. 80 cases (20.5%) had a preoperative EGD that showed HH and 39 of those (48.7%) had intraoperative findings consistent with HH. 301 (79%) patients had no HH on preoperative EGD, out of which 286 (95%) patients were also negative for HH intraoperatively. Compared with intraoperative diagnosis, EGD had sensitivity of 67% and specificity of 85%. The negative predictive value of EGD was 96% for HH but the positive predictive value was 34%.

Conclusions: EGD is a valuable informative tool in preoperative evaluation of bariatric patients. Our study demonstrated that EGD has high negative predictive value in ruling out HH but also with relatively low positive predictive value to prove its presence. Further studies are warranted to evaluate the discrepancies between the preoperative and intraoperative identification of HH, and standardization of definition for these findings.

Keywords

Bariatric surgery, Hiatal hernia, Sleeve gastrectomy, Esophagogastroduodenoscopy (EGD), Predictive value

Introduction

Obesity is a multinational public health burden as obesity has doubled since 1980 with 1.5 billion adults gone into obesity in 2008. In the United States, 34 percent of US adults 20-years-old and older are overweight, 34% obese, and 6% are Morbid obese [1]. Obesity is precisely linked to diabetes, hypertension and other chronic disease increase risk, Making big load on public health.

Preoperative upper gastrointestinal endoscopy (EGD) in patients going through bariatric surgery is argumentative [2]. It is supported routinely by some au-

thors to identify benign and cancerous pathology that largely leftovers without symptoms [2]. Others [3] advocate selective use, implying not much alter on surgical management of detected pathology, particularly in asymptomatic patients [4].

On the other hand, as (EGD) is an invasive procedure, other studies advocate that in preoperative assessment before bariatric surgery may be stand on the presence or absence of symptoms [5].

Nonetheless, many other studies [2] routinely performing EGD in preoperative bariatric surgery demonstrated that there is no correlation between manifestations and pathological findings on EGD. A variety of pathologies, including HH, esophagitis, gastritis, duodenitis, peptic ulcers, etc., can be determine on EGD [4]. Routine (EGD) preoperatively has not been constantly presented to change the management [1]. However, these endoscopic procedures contribute considerably to healthcare cost that may be avoided with selection and identification [6]. The purpose of this study is to de-

termine the value of preoperative endoscopy for bariatric surgery as a routine practice.

Material and Methods

Under IRB approval A retrospective review of 402 cases who underwent sleeve gastrectomy (SG) in our academic center between January 2011 and December 2015 was performed. We included all patients who had preoperative EDG. Twenty-one patients of revisional surgery after initial SG were excluded. We reviewed the EGD and operative reports for existence of hiatal hernia findings. Each patient’s preoperative EGD findings were compared with the intraoperative ones. We defined the

Table 3: HH repair (total no = 381).

Patients without repair		Patient with repair	
No	%	No	%
286	75%	31	25%

Table 4: Relation between endoscopic results and operative repair (total = 375).

Endoscopic results	Operative repair				X ²	P values
	Without repair		With repair			
	No	%	No	%	57.9	< 0.001
No HH	286	85.1	15	33.3		
HH	41	14.9	39	66.7		
Total	327	100	54	100		

Table 1: Patients with HH by EGD (total no = 381).

Patients without HH		Patient with HH	
No	%	No	%
301	79%	80	21%

HH: Hiatal Hernia.

Table 2: PPV & NPV.

	Sensitivity (95% CI)	Specificity (95% CI)	PPV (95% CI)	NPV (95% CI)	True positives	False positives	False negatives	True negatives
	%	%	%	%	NO.	NO.	NO.	NO.
Endoscopy	67% (50%-80%)	85% (81%-89%)	34% (24%-46%)	96% (92%-98%)	39	15	41	286

PPV: Positive Predictive Value; NPV: Negative Predictive Value.




Table 5: Other EGD Findings.

Findings:	No.	%
1- Barret Esophagus & HH	5	1.24%
2- Severe gastritis & HH	11	2.73%
3- Nodule in Duodenum	1	0.24%
4- Gastric Nodules	2	0.49%
5- Recurrent HH & Gastric fistula	1	0.24%

hiatal hernia by the EGD as a more than 2 cm separation of the caudally displaced esophago-gastric junction and diaphragmatic crural impression [7].

Statistical Analysis

Statistical presentation and analysis of the present study was conducted with SPSS V.18. and Data was expressed using X2 (*Chi 2*), Epi Cal 2000.

Results

From total of 402 SG cases, 381 (81% female) had preoperative EGD. The mean age of the subjects was 45.8 years (± 10.6 years) with the mean BMI of 47.5 kg/m² (± 8.8 kg/m²). There was no significant difference in age and BMI between males and females. 80 cases (20.5%) had a preoperative EGD that showed HH and 39 of those (48.7%) had intraoperative findings consistent with HH Table 1. 301 (79%) patients had no HH on preoperative EGD, out of which 286 (95%) patients were also negative for HH intraoperatively. Compared with intraoperative diagnosis, EGD had sensitivity of 67% ($\pm 7\%$) and specificity of 85% ($\pm 2\%$) Table 2. The negative predictive value of EGD was 96% ($\pm 2\%$) for HH but the positive predictive value was 34% ($\pm 5\%$) (Table 3 and Table 4) (Figure 1).

Discussion

Our study demonstrated that EGD has high negative predictive value in ruling out HH but also with relatively low positive predictive value to prove its presence.

As we already mentioned that (HH) and GERD is closely associated with morbid obesity. Also Sleeve gastrectomy can Lead to post-operative GERD symptoms [8].

In our practice we did not depend on the symptoms for the preoperative (EGD) we standardize it for all our patients, Zeni TM, et al. [9] and in Kuper MA, et al. [10] noticed that a high percentage of morbidly obese asymptomatic patients had positive findings During the screening. Only 12 of 69 patients (17.4%) had expressed upper gastrointestinal symptoms before the procedure. On the other hand, Heacock L, et al. (2012) [4] demonstrate that routine (EGD) produce a lot of endoscopic variations. In some cases, findings bring on a change in surgical management. But in most of the cases they did not make any change, Table 5 [11]. Furthermore, important endoscopic findings are common among obese patients which may alter the surgical plan or even abort it. While some authors founded the negative predictive value in low-risk patients was not sufficient, finally they

recommend routine (EGD) for all patients Lee J, et al. [8].

Per De Palma, G. D. and P. Forestieri [12]. They mentioned that in some selective procedures changes may be done if meaningful upper GI findings present, like large HH or Barrett's esophagus, and was detected preoperatively by EGD. The published guidelines of the European Association for Endoscopic Surgery state that esophagogastroduodenoscopy (EGD), or upper-GI series, is advisable for all bariatric procedures. Similarly, the recently published guidelines from the ASGE recommended that EGDS should be performed in all symptomatic patients undergoing bariatric surgery. A shortage of interconnection between patient manifestations and EGD Finding has been discussed by many studies, implying that routine preoperative endoscopy can be useful in HH diagnosis [12]. Although, putting in consideration the relatively less important finding in the majority of lesions discovered on routine EGDS, cost and the amount of secondary irrelevant workup, several studies have instead suggested a non-endoscopic approach for patients without symptoms [12].

One of the major burden of performing routine preoperative upper endoscopy is the hazard of sedation. Cardiopulmonary complications are the worst complications accompanying the sedation during the procedure, with a mortality rate of 0.03% and a serious morbidity rate of 0.54% [13]. In morbidly obese patients, the complications will be even higher because the possibility of restrictive lung disease, obstructive sleep apnea, pulmonary hypertension, and cardiac diseases. It is of cardinal importance that procedures in obese patients are done in a fully supplied setting and by a strong-organized team endoscopists and anesthesiologists Sharma, et al. [14]. However, finding, a single cancer may not be cost effective, there are obvious benefits to that one patient, also strict to a medico-legal standard of care InMong C, et al. [15].

On the other hand, Barium studies and other contrast imaging may be another option for diagnose HH, however, the advantages of the EGD goes higher the diagnosis only but the confirmation of the pathology and findings by biopsy is achievable. Furthermore, the finding during the operation still has the most accurate evidence. In the study of Broucek JR, et al. [16] showed that UGI series have poor positive and negative predictive values in preoperatively diagnosing HH. Also, patient symptoms were different subjectively and anti-reflux medication did not match with radiologic or intraoperative findings of HH [17]. Standardizing routine preoperative EGD before bariatric surgery Still a topic for arguments. Although small HH are over estimated with EGD, and Endoscopist experiences also has a big role Mohammed R, et al. [18]. However, due to high negative predictive value of the EGD in ruling out hiatal hernia, we can avoid hiatal dissection in these patients who did not have evidence of HH on the Preoperative EGD.

Conclusions

EGD is a valuable informative tool in preoperative evaluation of bariatric patients. Our study demonstrated that EGD has high negative predictive value in ruling out HH but relatively low positive predictive value to prove its presence. Further studies are warranted to evaluate the discrepancies between the Preoperative and intraoperative identification of HH and standardization of definition of operative finding of hiatal hernia in this group of patients undergoing sleeve gastrectomy.

Conflict of Interest

The authors declare that they have no conflict of interest.

Ethical Statement

All procedures involving human participants were performed in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki Declaration and its later amendments or comparable ethical standards.

Informed consent was obtained from all individual participants included in the study.

Authors have no pertinent disclosures.

References

- Sharaf RN, Weinshel EH, Bini EJ, Rosenberg J, Sherman A, et al. (2004) Endoscopy Plays an Important Preoperative Role in Bariatric Surgery. *Obesity Surgery* 14: 1367-1372.
- de Moura Almeida A, Cotrim HP, Santos AS, Bitencourt AG, Barbosa DB, et al. (2008) Preoperative upper gastrointestinal endoscopy in obese patients undergoing bariatric surgery: is it necessary? *Surg Obes Relat Dis* 4: 144-149.
- Asge Standards of Practice Committee, Anderson MA, Gan SI, Fanelli RD, Baron TH, et al. (2008) Role of endoscopy in the bariatric surgery patient. *Gastrointest Endosc* 68: 1-10.
- Praveenraj P, Gomes RM, Kumar S, Senthilnathan P, Parathasarathi R, et al. (2015) Diagnostic Yield and Clinical Implications of Preoperative Upper Gastrointestinal Endoscopy in Morbidly Obese Patients Undergoing Bariatric Surgery. *J Laparoendosc Adv Surg Tech A* 25: 465-469.
- Abd Ellatif ME, Alfalah H, Asker WA, El Nakeeb AE, Magdy A, et al. (2016) Place of upper endoscopy before and after bariatric surgery: A multicenter experience with 3219 patients. *World J Gastrointest Endosc* 8: 409-417.
- Gómez V, Bhalla R, Heckman MG, Florit PT, Diehl NN, et al. (2014) Routine Screening Endoscopy before Bariatric Surgery: Is It Necessary? *Bariatric Surg Pract Patient Care* 9: 143-149.
- Van Weyenberg SJB (2013) Diagnosis and Grading of Sliding Hiatal Hernia. *Video Journal and Encyclopedia of GI Endoscopy* 1: 117-119.
- Lee J Wong SK, Liu SY, Ng EK (2017) Is Preoperative Upper Gastrointestinal Endoscopy in Obese Patients Undergoing Bariatric Surgery Mandatory? An Asian Perspective. *Obes Surg* 27: 44-50.
- Zeni TM, Frantzides CT, Mahr C, Denham EW, Meiselman M, et al. (2006) Value of preoperative upper endoscopy in patients undergoing laparoscopic gastric bypass. *Obes Surg* 16: 142-146.
- Küper MA, Kratt T, Kramer KM, Zdichavsky M, Schneider JH, et al. (2010) Effort, safety, and findings of routine preoperative endoscopic evaluation of morbidly obese patients undergoing bariatric surgery. *Surg Endosc* 24: 1996-2001.
- Soricelli E, Iossa A, Casella G, Abbatini F, Cali B, et al. (2013) Sleeve gastrectomy and crural repair in obese patients with gastroesophageal reflux disease and/or hiatal hernia. *Surg Obes Relat Dis* 9: 356-361.
- De Palma GD, Forestieri P (2014) Role of endoscopy in the bariatric surgery of patients. *World J Gastroenterol* 20: 7777-7784.
- Zakeri N, Coda S, Webster S, Howson W, Thillainayagam AV (2015) Risk factors for endoscopic sedation reversal events: a five-year retrospective study. *Frontline Gastroenterol* 6: 270-277.
- Sharma VK, Nguyen CC, Crowell MD, Lieberman DA, de Garmo P, et al. (2007) A national study of cardiopulmonary unplanned events after GI endoscopy. *Gastrointest Endosc* 66: 27-34.
- Mong C, Van Dam J, Morton J, Gerson L, Curet M, et al. (2008) Preoperative endoscopic screening for laparoscopic Roux-en-Y gastric bypass has a low yield for anatomic findings. *Obes Surg* 18: 1067-1073.
- Broucek JR, Ritter LA, Francescatti AB, Smith CH, Luu MB, et al. (2014) Radiographic predictability of hiatal hernia prior to gastric band surgery. *JLS* 18: 243-245.
- Fornari F, Gurski RR, Navarini D, Thiesen V, Mestriner LH, et al. (2010) Clinical utility of endoscopy and barium swallow X-ray in the diagnosis of sliding hiatal hernia in morbidly obese patients: a study before and after gastric bypass. *Obes Surg* 20: 702-708.
- Mohammed R, Fei P, Phu J, Asai M, Antanavicius G (2017) Efficiency of preoperative esophagogastroduodenoscopy in identifying operable hiatal hernia for bariatric surgery patients. *Surg Obes Relat Dis* 13: 287-229.