



CASE REPORT

A Novel Mutation in the MFN2 Gene Associated with Hereditary Sensory and Motor Neuropathy with Proximal Predominance (HMSN-P)- A Case Report

D Hettiarachchi*, T K Wethasinghe, N F Neththikumara, BAPS Pathirana and VHW Dissanayake

Human Genetics Unit, Faculty of Medicine, University of Colombo, Sri Lanka

*Corresponding author: Dr. Dineshani Hettiarachchi, Lecturer, Department of Anatomy, Faculty of Medicine, University of Colombo, Sri Lanka, Tel: +9477-722-2228



Abstract

Background: Mutations in mitofusin 2 gene have been reported in Charcot-Marie-Tooth type 2 disease also known as Hereditary Sensory and Motor Neuropathy. With its cytogenetic location: 1p36.22.

Case presentation: A 43-year-old female with a family history of neuropathy was experiencing gradual deterioration and proximal weakness of the bilateral lower limb for the past 3 years. Her MRI scan (Brain and whole Spinal) was normal and Electromyography (EMG) report was suggestive of motor & sensory demyelinating polyneuropathy with features of segmental involvement. Her Creatine phosphokinase (CPK) levels were marginally elevated. However, all other investigations were within normal range.

Conclusions: A heterozygous missense variant denoted as c.701T > A at level of cDNA in exon 7 of the MFN2 gene was discovered which resulted in the substitution of Methionine by Lysine at position 234 [*p.Met234Lys*] of the amino acid sequence which was confirmed by Sanger sequence.

Keywords

Hereditary motor and sensory neuropathies, Exome sequencing, MFN2 gene

Background

Hereditary motor and sensory neuropathy with proximal dominance (HMSN-P) has been reported as a rare type of autosomal dominant adult-onset Charcot-Marie-Tooth disease. Hereditary motor sensory neuropathies (HMSN) are a group of heterogeneous diseases classified under motor neuropathies [1]. In Charcot-

Marie-Tooth disease (CMT) and the hereditary sensory neuropathies there is a significant sensory involvement along with distal lower motor neuron weakness [2]. The classical phenotype of HMSN is a length-dependent motor weakness and atrophy, primarily affecting the intrinsic foot muscles and the peroneal compartment of the leg, often leading to foot deformities such as pes cavus, pes planus, and clawing of the toes [3].

Mutations in the mitofusin 2 (MFN2) gene, which encodes a mitochondrial GTPase mitofusin protein, have recently been reported to cause both Charcot-Marie-Tooth 2A (CMT2A) and hereditary motor and sensory neuropathy VI (HMSN VI) [4].

Case Presentation

A 43-year-old female with a family history of neuropathy presented with bilateral progressive proximal muscle weakness with sensory involvement. She first noticed reduced sensation and muscle weakness 3 years back and was clinically diagnosed to have hereditary sensory motor neuropathy. Her MRI scan (Brain and whole Spinal) was normal nerve conduction studies (NCS) and Electromyography (EMG) report was suggestive of motor & sensory demyelinating polyneuropathy with features of focal segmental involvement. Her creatinine phosphokinase (CPK) levels were elevated. However, all other investigations were within normal range. Proband's father who had died at the age of 54 yrs. due to an unknown cause had symptoms of sensory

motor neuropathy, which manifested at the age of 24 years. Proband's two daughters age 14 and 17 years are asymptomatic and was found to be free of the mutation (Figure 1). Whole exome sequencing of the proband revealed a previously unreported heterozygous missense Variant in the *MFN2* gene which is likely pathogenic.

Whole exome sequencing

Whole exome sequencing (WES) of the proband's DNA was performed in line with the SureSelect® Human All Exon V6 kit on an Illumina® HiSeq 4000 Next Generation Sequencer. The DNA sequence was compared with the UCSC hg19 reference sequence. All reportable sequence variants were confirmed by visual inspection of the alignment.

Bioinformatics analysis

The sequence reads were analyzed using in-house variant calling and annotation pipeline. Reads were aligned to a human reference sequence and processed

using standard software. The data obtained were analyzed by annotation with multiple INFO fields required for filtering

(dbSNP allele frequencies (<http://www.ncbi.nlm.nih.gov/projects/SNP/>),

1000 Genomes allele frequencies (www.1000genomes.org),

Exome Aggregation Consortium (ExAC) allele frequencies (http://exac.broadinstitute.org),

the OMIM database and the Clinvar database (<https://www.ncbi.nlm.nih.gov/clinvar/>), using the SnpSift tool. Variant effect prediction was then performed using Genetic variant annotation and the effect prediction toolbox (SnpEff). Sanger sequencing was done to confirm the presence of the variants of interest by analyzing the chromatogram using Codon Code Aligner software.

In silico analysis

In silico functional prediction was performed using Mutation Taster, SIFT, Polyphen2, and Provean.

A heterozygous missense mutation denoted as *c.701T > A* at level of cDNA in exon 7 of the *MFN2* gene was discovered which resulted in the substitution of Methionine by Lysine at position 234 [*p.Met234Lys*] of the amino acid sequence was confirmed by Sanger sequence (Figure 2). Substitution of methionine to Lysine changes the physio-chemical characteristics of the amino acid, thus causes a non-conservative amino acid substitution.

This substitution in a highly conserved region of the *MFN2* gene (phastCons100way Vertebrate score = 1; PhyloP score = 5.145) and majority of in-silico functional prediction tools concluded this variant as pathogenic (Table 1).

Discussion and Conclusions

Charcot-Marie-Tooth disease (CMT), also named hereditary motor and sensory neuropathies, includes a clinically and genetically heterogeneous group of disorders affecting the peripheral nervous system, with a relatively high prevalence of 1:2,500 [5]. The main forms of




Figure 1: Pedigree of the proband's family with affected individuals.

Table 1: Results of in silico mutation prediction analysis.


	Prediction	Score
Provean a	Deleterious	-4.33
SIFT b	Damaging	0.001
Mutation Taster c	Disease causing	0.999
PolyPhen2 d	Benign (HumDiv- model)	0.007
PolyPhen2 d	Benign (HumVar- model)	0.047

a) <http://provean.jcvi.org/index.php> ; score threshold is -2.5 for binary classification.

b) http://sift.jcvi.org/www/SIFT_chr_coords_submit.html threshold < 0.05.

c) <http://www.mutationtaster.org>; Scores range from 0.0 to 1.0.

d) <http://genetics.bwh.harvard.edu/pph2>; deleterious threshold > 0.5.



CMT have been subdivided into primary demyelinating neuropathies (myelinopathies), which include CMT1 (autosomal-dominant inheritance), CMT1X (X-linked), and CMT4 (autosomal-recessive), and neuropathies characterized by primary axonal defects such as CMT2 (autosomal-dominant or recessive). Additional sub-forms of hypomyelinating and/or dysmyelinating neuropathies include Dejerine-Sottas syndrome (DSS, also called CMT3), hereditary neuropathies with liability to pressure palsies (HNPP), and congenital hypomyelination (CH) [6]. Mutations in mitofusin 2 (*MFN2*) have been reported in Charcot-Marie-Tooth type 2 (CMT2) families. In patients with a documented family history of CMT2 the frequency of *MFN2* mutations was 33% indicating that *MFN2* mutations are a major cause in this population [7].

As with our patient most suffer from demyelinating forms of CMT.

However, investigations of the underlying molecular and cellular disease mechanisms, employing cell cultures and animal models have revealed an overlap of pathological signs between myelinopathies and axonopathies [8] Charcot-Marie-Tooth disease type 2A was previously mapped to an interval containing *KIF1B*. New studies have shown that CMT2A patients contain a loss-of-function mutation in the motor domain of the *KIF1B* gene. This is clear indication that defects in axonal transport due to a mutated motor protein can underlie human peripheral neuropathy [9]. However, the gene responsible

for CMT2A is still controversial, and the mechanism of pathogenesis remains uncertain. It has also been shown that the receptor tyrosine kinase *IGF1R* is a new direct binding partner of *KIF1Bβ*, and its binding and transport is specially impaired by the Y1087C mutation of *KIF1Bβ*, which we detected in hereditary neuropathic patients [10]. Approximately ten genes have now been identified to cause CMT disease [11]. Whole exome sequencing (WES) of the proband revealed a novel variant in the *MFN2* gene, c.701T > A (p.Met234Lys).

MFN2 is a mitochondrial transmembrane GTPase, which regulates mitochondrial architecture by fusion of mitochondria and most pathological variants of this gene is located within or immediately upstream of the GTPase domain or within two coiled-coil domains, which are critical for the functioning or mitochondrial targeting of *MFN2*. Formation of a mitochondrial network is essential for a proper functioning PNS [11]. Even though there is no current treatment for the condition recent studies have shown that antisense oligonucleotides (ASOs) offer new hope for patients with Charcot-Marie-Tooth disease type 1A (CMT1A) [12]. In conclusion we report a novel heterozygous missense variant in exon 7 of the *MFN2* gene causing CMT disease.

Declarations

Ethics approval and consent to participate

Ethics approval for the study was obtained by the

Faculty of Medicine, University of Colombo Ethics Review Committee (EC-16-179). Duly filled consent forms are available with the corresponding author.

Consent to publish

Written informed consent to publish this material was obtained from the proband and her two daughters.

Availability of data and materials

Data is not available in a public domain it will be shared on request from interested parties.

Competing interests

Authors declare that there are no competing interests.

Funding

There was no funding for this study.

Authors' contributions

DH and VHWD were the clinicians looking after the patient. BAPSP and KW performed laboratory testing. NN performed bioinformatics analysis. DH wrote the first draft of the manuscript with contributions from all. All authors reviewed, modified and approved the final version of the manuscript.

Acknowledgements

We wish to acknowledge the referring doctor and the laboratory staff for their contribution.

References

1. Lee SS, Lee HJ, Park JM, Hong YB, Park KD, et al. (2013) Proximal dominant hereditary motor and sensory neuropathy with proximal dominance association with mutation in the TRK-fused gene. *JAMA Neurol* 70: 607-615.
2. Rossor AM, Kalmar B, Greensmith L, Reilly MM (2012) The distal hereditary motor neuropathies. *J Neurol Neurosurg Psychiatry* 83: 6-14.
3. Dyck PJ, Litchy WJ, Minnerath S, Bird TD, Chance PF, et al. (1994) Hereditary motor and sensory neuropathy with diaphragm and vocal cord paresis. *Ann Neurol* 35: 608-615.
4. Chung KW, Kim SB, Park KD, Choi KG, Lee JH, et al. (2006) Early onset severe and late-onset mild Charcot-Marie-Tooth disease with mitofusin 2 (MFN2) mutations. *Brain* 129: 2103-2118.
5. Skre H (1974) Genetic and clinical aspects of Charcot-Marie-Tooth's disease. *Clin Genet* 6: 98-118.
6. Berger P, Young P, Suter U (2002) Molecular cell biology of Charcot-Marie-Tooth disease. *Neurogenetics* 4: 1-15.
7. Verhoeven K, Claeys KG, Züchner S, Schröder JM, Weis J, et al. (2006) MFN2 mutation distribution and genotype/phenotype correlation in Charcot-Marie-Tooth type 2. *Brain* 129: 2093-2102.
8. Eva Nelis, Neva Haites, Christine Van Broeckhoven (1999) Mutations in the peripheral myelin genes and associated genes in inherited peripheral neuropathies. *Human Mutation* 13: 11-28.
9. Zhao C, Takita J, Tanaka Y, Setou M, Nakagawa T, et al. (2001) Charcot-Marie-Tooth disease type 2A caused by mutation in a microtubule motor KIF1B β . *Cell* 105: 587-597.
10. Xu F, Takahashi H, Tanaka Y, Ichinose S, Niwa S, et al. (2018) KIF1B β mutations detected in hereditary neuropathy impair IGF1R transport and axon growth. *J Cell Biol* 217: 3480-3496.
11. Kijima K, Numakura C, Izumino H, Umetsu K, Nezu A, et al. (2005) Mitochondrial GTPase mitofusin 2 mutation in Charcot-Marie-Tooth neuropathy type 2A. *Hum Genet* 116: 23-27.
12. Zhao HT, Damle S, Ikeda-Lee K, Kuntz S, Li J, et al. (2018) PMP22 antisense oligonucleotides reverse Charcot-Marie-Tooth disease type 1A features in rodent models. *J Clin Invest* 128: 359-368.