



CASE REPORT

Floating First Metacarpal: Fracture Dislocation of the Trapezio-Metacarpal Joint Associated with Fracture of the Base of the Proximal Phalanx: A Case Report

Hamdi Kaziz*, Thabet Mouelhi, Walid Osman, Nader Naouar and Mohamed Laziz Ben Ayeche

Department of Orthopedics, Hospital Sousse, Tunisia

*Corresponding author: Hamdi Kaziz, Department of Orthopedics, Hospital Sousse, Tunisia, Tel: +21698366196, E-mail: hamdi.kaziz@gmail.com

Abstract

Floating first metacarpal has been scarcely described in the literature. Fracture dislocation of the trapezio-metacarpal joint has been reported more than the isolated form but deemed exceptional. We report a 40-year-old man consulting for an injury on his right thumb in which the described mechanism was an axial metacarpal force with both flexed trapezio-metacarpal and interphalangeal joints. X-ray findings revealed a floating metacarpal composed from a Bennett fracture dislocation of the trapezio-metacarpal joint and a fracture dislocation of the base of the proximal phalanx. A closed reduction and percutaneous fixation was achieved. After removal of pins and physiotherapy, and at 12 months of follow up, anatomical and functional outcomes were satisfied with a normal motion of the thumb. The aim of our case report is to improve the closed reduction and percutaneous fixation on fracture dislocation of the trapezio-metacarpal on contrast to the isolated dislocation form in which closed reduction must be avoided.

Keywords

Trapezio-metacarpal, Fracture, Dislocation, Bennet, Phalanx, Floating, Closed reduction, Fixation

Introduction

Traumatic trapezio-metacarpal joint fracture dislocation is exceptional [1]. Several treatment options have been described; however, optimal treatment strategies are controversial [2]. In this article, we report a case of trapezio-metacarpal joint fracture dislocations composed from Bennett fracture and horizontal disloca-

tion of the joint associated with a fracture of the base of the proximal phalanx of the thumb. The aim of our case report is to put into consideration the main characteristics of such entity and review the different options of treatment. The fracture dislocation was successfully treated with closed reduction and percutaneous K-wires fixation. To our knowledge, this association has never been scarcely reported reviewing the literature [3,4]. The functional and anatomical outcomes revealed that closed reduction and percutaneous fixation is available in fracture dislocation more than isolated dislocation.

Case Report

We report a 40-year-old man, consulting in emergency department after a traffic road accident, complained about pain, tenderness, and swelling of his right thumb. X-ray findings revealed a horizontal dislocation of the trapezio-metacarpal joint combined to a Bennett fracture type III and a fracture of the base of the proximal phalanx of the same thumb (Figure 1). The diagnosis of floating first metacarpal is so taken. Under general anesthesia, a closed reduction of trapezio-metacarpal joint fracture dislocation and the phalange fracture was achieved under fluoroscopic control. Repeated X-ray revealed a perfect reduction of both dislocation and fractures without any incongruence in both of the two joints. The reduced dislocation was fixed by percutaneous pinning according to Iselin procedure whereas; the phalange fracture was fixed by one direct k-wire (Figure 2). The thumb was immobilized with cast for 6 weeks.




Figure 1: (A,B) X-ray views of the right hand revealing the floating metacarpal of the thumb with a fracture dislocation of the trapezio-metacarpal joint composed from an horizontal dislocation with bennet fracture and a fracture of the base of the proximal phalanx.




Figure 2: (A,B) Post operative X-ray views: reduction of the dislocation and fixation according to Iselin procedure.

Physiotherapy and rehabilitation of the thumb were started after the removal of the cast and maintained for two months. At the 12-month of follow-up, physical examination showed an unrestricted painless range of motion of the thumb: palmar abduction 70°, radial abduction 50°, complete and symmetrical opposition and retraction (Figure 3). Grip strength was 37 kg against 40 kg on the uninjured right side; The pinch power was 7 kg on both sides. An excellent radiological result (Figure 4) and functional outcome was noted with the patient performing normally at his job, having no limitation of his activities of daily living, and reporting no pain or instability symptoms.

Discussion

The trapezio-metacarpal fracture-dislocation of the thumb is exceptional. It accounts for less than 1% of all hand injuries [1]. The carpometacarpal joint of the thumb is a biconcave saddle joint. The trapezium is convex on anteroposterior views and concave on lateral

views, however, the thumb metacarpal has the opposite views. In fact, this typical anatomical configuration allows the thumb to have a large range of motion varying from abduction to opposition while the joint remains stable [2]. Besides, the trapezio-metacarpal joint is stabilized by a thickened joint capsule composed of several ligaments but stability is primarily provided by 4 main ligaments: the volar oblique, the intermetacarpal, the dorsoradial, and the posterior oblique ligament [5]. Despite the variety of biomechanical studies, no ligament has been identified to be the key stabilizer of the joint so that no ligament is identified to be the main damaged one in each kind of dislocation [6]. However, and for a long period of time, the volar oblique ligament has been believed to be the key stabilizer of the trapezio-metacarpal joint [7]. This intrinsic joint stability makes isolated trapezio-metacarpal dislocation quite rare injury. So, the fracture dislocation occurs more than the isolated form [8]. There are evidences that, in trapezio-metacarpal dislocation, the dorsoradial ligament tear, lead




Figure 3: Functional outcome of the thumb. Motion of the trapezio-metacarpal joint and interphalangeal joint seems to be similar to the other one (A) Extension of the thumb in dorsal view; (B) Extension of the thumb in volar view; (C) Flexion of the thumb; (D) Dorsal view of thumb opposition; (E) Volar view of thumb opposition; (F) Dorsal view of the first commissural spaces of the hands.



Figure 4: (A,B) X-ray views at the final follow up showing joint congruency of the trapezio-metacarpal and interphalangeal joint.

to a subperiosteally stripping of the volar oblique ligament deemed intact. Therefore, extension and prona-

tion of the thumb tightens the stripped volar oblique ligament and caused dislocation of the joint associated

with a partial fracture of the metacarpal base [6-9]. In fact, this anatomic pathology reveals that healing the dorsolateral ligament in an appropriate position, the trapezio-metacarpal joint becomes stable [10]. Several studies have reported some mechanisms to explain this injury but two mechanisms have been adopted for a traumatic fracture-dislocation of the trapezio-metacarpal [11,12]. The first one is a longitudinally directed force along the axis of the metacarpal with a full flexion of the joint [11,12]. The second is a force driven into the first web space of the thumb. This force separates the base of the first and second metacarpal joints and produces a fracture dislocation of the trapezio-metacarpal joint [11,12]. In our case, longitudinally directed force along the axis of the metacarpal with flexion of both trapezometacarpal and interphalangeal joint is the mechanism of floating metacarpal. The treatment of this injury is controversial [13]. Several options have been reported in literature [13]. The treatment's procedures varied from a closed reduction and immobilization to a closed or open reduction and K-wire fixation with or without the reconstruction of the capsule and ligaments [14]. It is well reported that the objective of the management of acute trapezio-metacarpal fracture-dislocations is to ensure the joint stability after reduction [15]. Open reduction is indicated after failure of the closed reduction.

We performed a closed reduction and percutaneous K-wires fixation, according to Iselin procedure [16] for the trapezio-metacarpal fracture dislocation and a direct pinning of the base in the phalange fracture. Although the extent of ligamentous injury was not determined, we believed that the volar oblique ligament was intact since the avulsion of the first metacarpal base was seen. Open reduction and ligamentous repair were discussed for only the isolated dislocation of the joint since there is a total disruption of the dorsal capsule and the dorsoradial ligament [13,14,17]. Whereas, in our case, the metacarpal base and the trapezium bone remains connected by these structures. Hence, Ligament reconstruction of the inter metacarpal and capsular structures, such as an inter metacarpal abductor pollicis longus ligamentoplasty [18] which have been recommended following isolated dislocation in order to prevent post traumatic arthritis and decrease the progression of the degenerative process seem no mandatory in fracture dislocation of the joint and deemed only available in isolated dislocation [19]. In our case report, and at the 12-months of follow-up, the patient showed a normal range of motion, normal strength and no joint instability. Overall, he was satisfied with the outcome. The aim of our case report is to remember that closed reduction and percutaneous pinning on fracture dislocation in trapezio-metacarpal joint deemed an available treatment option. Reparation of ligament and capsular damages is relevant step since the radio dorsal ligament and dorsal capsule are intact.

Conclusion

Floating metacarpal thumb has been scarcely described to our knowledge. The mechanism of injury is similar to the isolated form but with a flexion of the interphalangeal joint of the thumb. If these injuries are not accurately diagnosed and treated, obviously, they cause pain and limitation of motion due to arthritis. Reconstruction of ligament and capsular damages may not be necessary in trapezio-metacarpal fracture-dislocations in where the metacarpal base fragment and the trapezium remain connected by the dorsal structures. A percutaneous stabilization after closed reduction will ensure a good outcome.

Acknowledgement

I would like to express my deepest appreciation to all those who have been contributed to the achievement of this case report.

References

- Mueller JJ (1986) Carpometacarpal dislocations: report of five cases and review of the literature. *J Hand Surg Am* 11: 184-188.
- Bosmans B, MHJ Verhofstad, T Gosens (2008) Traumatic thumb carpometacarpal joint dislocations. *The Journal of Hand Surgery* 33: 438-441.
- Messaoudi T, Errhaimini M, Ghoubach M, Chafik R, Madhar M, et al. (2015) Floating thumb metacarpal in a motorcyclist: A case report. *Chir Main* 34: 91-93.
- Rajeev A, Noureldin S, Graham D (2014) Floating thumb with double dislocation of carpometacarpal and metacarpophalangeal joints. *International Journal of Surgery Case Reports* 5: 1102-1105.
- Bettinger PC, Linscheid RL, Berger RA, Cooney WP III, Kainan A, et al. (1999) An anatomic study of the stabilizing ligaments of the trapezium and trapeziometacarpal joint. *The Journal of Hand Surgery* 24: 786-798.
- Strauch RJ, Behrman MJ, Rosenwasser MP (1994) Acute dislocation of the carpometacarpal joint of the thumb: an anatomic and cadaver study. *J Hand Surg Am* 19: 93-98.
- Eaton RG, Littler JW (1973) Ligament reconstruction for the painful thumb carpometacarpal joint. *J Bone Joint Surg* 55: 1655-1666.
- Henry M (2001) Fractures and dislocations of the hand. In: Bucholz RW, Heckman JD, Rockwood, Green's fractures in adults. (5th edn), Lippincott, Philadelphia, 655-748.
- Wee JT, Chandra D, Satku K (1988) Simultaneous dislocations of the interphalangeal and carpometacarpal joints of the thumb: a case report. *J Hand Surg* 13: 224-226.
- Ahmadreza Afshar, Fardin Mirzatoeloei (2005) Carpometacarpal joint dislocation of the thumb associated with the fracture of the trapezium. *Archives of Iranian Medicine* 9: 282-283.
- Johnson SR, Jones DG, Hoddinott HC (1987) Missed carpometacarpal dislocation of the thumb in motorcyclists. *Injury* 18: 415-416.
- Shah J, Patel M (1983) Dislocation of the carpometacarpal joint of the thumb: a report of four cases. *Clin Orthop Relat Res* 166-169.

13. Changhoon Jeong, Hyoung-Min Kim, Sang-Uk Lee, Il-Jung Park (2012) Bilateral Carpometacarpal Joint Dislocations. *Clin Orthop Surg* 4: 246-248.
14. Simonian PT, Trumble TE (1996) Traumatic dislocation of the thumb carpometacarpal joint: early ligamentous reconstruction versus closed reduction and pinning. *J Hand Surg Am* 21: 802-806.
15. Watt N, Hooper G (1987) Dislocation of the trapezio-metacarpal joint. *J Hand Surg Br* 12: 242-245.
16. M Iselin (1960) Atlas de technique opératoire. Chirurgie de la main. *JBJS Am* 42: 1453-1492.
17. Razemon JP (1984) Carpal bones fractures. In: Tubiana R, *Treaty of hand surgery*. Tome II. Paris: Masson, 667-692.
18. Rocchi L, Merolli A, Cotroneo C, Morini A, Brunelli F, et al. (2011) Abductor pollicis longus hemitendon looping around the first intermetacarpal ligament as interposition following trapeziectomy: a one-year follow-up study. *Orthop Traumatol Surg Res* 97: 726-733.
19. Kukreti S, Harrington P (2004) Carpometacarpal joint dislocation of the thumb associated with fracture of the trapezium: a case report. *European Journal Of Orthopaedic Surgery & Traumatology* 14: 38-39.